

Association of vitamin D receptor gene polymorphisms in Chinese patients with generalized aggressive periodontitis

S. Li¹, M. H. Yang², C. A. Zeng¹,
W. L. Wu², X. F. Huang², Y. Ji²,
J. Q. Zeng¹

¹Department of Stomatology, Medical School, Jianggangshan University, Ji'an, Jiangxi, China and ²Affiliated Stomatology Hospital, Medical School, Nanjing University, Nanjing, Jiangsu, China

Li S, Yang MH, Zeng CA, Wu WL, Huang XF, Ji Y, Zeng JQ. Association of vitamin D receptor gene polymorphisms in Chinese patients with generalized aggressive periodontitis. *J Periodont Res* 2008; 43: 360–363. © 2008 The Authors. Journal compilation © 2008 Blackwell Munksgaard

Background and Objective: The clinical features suggest that genetic factors may have a strong influence on susceptibility to aggressive periodontitis. The aim of this study was to investigate the association of vitamin D receptor gene polymorphisms with generalized aggressive periodontitis in Chinese patients.

Material and Methods: A restriction fragment length polymorphism (RFLP) for 10,438,141 C to T (rs1544410, *BsmI*), 10,382,063 A to G (rs731236, *TaqI*), 10,382,143 C to A (rs7975232, *ApaI*) and 10,416,201 A to G (rs2228570, *FokI*) of vitamin D receptor gene was analysed by polymerase chain reaction, followed by digestion with restriction enzymes and gel electrophoresis. The genotypes of 51 generalized aggressive periodontitis patients and 53 periodontally healthy control subjects were analysed. The genotypic and allelic frequencies of each polymorphism site for the patients and control subjects were compared.

Results: The distribution of vitamin D receptor *FokI* genotypes and alleles between the two groups was significantly different ($p = 0.043$ and $p = 0.012$, respectively). The F allele seemed to increase the susceptibility of aggressive periodontitis (odds ratio = 2.02, 95% confidence interval = 1.16–3.50) in Chinese patients. There was no significant difference in the genotype distribution or the allele frequencies of vitamin D receptor *BsmI*, *ApaI* and *TaqI* between two groups.

Conclusion: The study indicates that *FokI* polymorphism of vitamin D receptor gene might be associated with generalized aggressive periodontitis in Chinese patients. In addition, the carriage of F allele increases the risk of developing generalized aggressive periodontitis.

Ming Hua Yang, Affiliated Stomatology Hospital, Medical School, Nanjing University, 30 ZhongYang Road, Nanjing, Jiangsu 210008, China
Tel: +86 025 83620252
Fax: +86 025 83620202
e-mail: y_mh@hotmail.com

Key words: vitamin D receptor; aggressive periodontitis; polymorphism; susceptibility

Accepted for publication August 6, 2007

Generalized aggressive periodontitis (GAgP) represents a group of inflammatory diseases with common pathogenic features, such as onset primarily during early adult years and generalized rapid destruction of periodontal tissue, leading to loss of teeth. The

aetiology of GAgP involves complex interactions between exogenous factors, in particular periodontal bacteria, and genetic factors (1). Although periodontopathogenic bacteria may indeed constitute an initiator of the condition, its clinical features have indicated that

genetic factors may have a strong influence on susceptibility to GAgP.

Vitamin D has special roles in regulating the metabolism of calcium and phosphorus, and in the immune system. The vitamin D receptor (VDR) is involved in a variety of biological

processes, including bone metabolism and the modulation of the immune response. It is clear that mutations in functionally critical areas of the VDR gene can have profound effects on mineral metabolism and bone mineral density (2,3). Vitamin D receptor is also an important nuclear receptor of vitamin D₃, which is known to be essential for the maintenance of mineral homeostasis and for bone structural integrity. Vitamin D₃ and its receptor have also been demonstrated to act as upregulating agents during innate immunity (4). Vitamin D receptor mediates these effects via a variety of mechanisms, including transcription regulation, mRNA stability and post-translational modifications.

The aim of this study was to determine the prevalence of *BsmI*, *ApaI*, *TaqI* and *FokI* VDR genotypes in Chinese GAgP patients and to evaluate the association of VDR gene polymorphisms with GAgP.

Materials and methods

Subjects

A total of 51 patients with GAgP were recruited from the periodontic clinics at the Affiliated Stomatology Hospital, Nanjing University. The GAgP patient group contained 28 males and 23 females, with an age range from 16 to 49 years (mean 35.7 years). Generalized aggressive periodontitis was diagnosed on the basis of clinical attachment loss and radiographic patterns of alveolar bone loss, using the criteria defined by the American Academy of Periodontology (5). The main clinical criteria of GAgP patients recruited in this study were a clinical attachment loss of at least 6 mm or alveolar bone resorption of one-third of the length of the root, affecting at least 10 teeth, four or more of which were not incisors or first molars. In addition, 53 healthy control subjects with no previous or existing periodontal disease were selected from the volunteers. The healthy control subjects ranged in age from 24 to 45 years (mean 39.1 years). All subjects were Chinese Han nationality, and subjects with a history of systemic disease or

smoking were excluded from the study. The nature of the study was explained to all subjects verbally and in writing and all signed a content form.

Sample collection and DNA extraction

Buccal swab samples were taken from each subject. The DNA was extracted from the buccal swab samples with a buccal swab DNA extraction reagent (Tiangen, Beijing, China), following the manufacturer's instructions. The concentration of each DNA sample was determined by ultraviolet spectrophotometry.

Genotype analysis

Four single-nucleotide polymorphisms (SNPs) of the VDR gene were analysed in this study, at the following positions (according to GenBank accession no. NT_029419): 10,438,141 C to T (rs1544410, *BsmI*), 10,382,063 A to G (rs731236, *TaqI*), 10,382,143 C to A (rs7975232, *ApaI*) and 10,416,201 A to G (rs2228570, *FokI*). The four sites were analysed by the polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP). The PCR reaction primers were designed by Primer Premier 5 (Premier Biosoft International, Palo Alto, CA, USA). All primers and their amplifying fragment length are shown in Table 1.

Reaction conditions and cycling parameters were as follows: 200 ng of genomic DNA was used for PCR amplification in a reaction mixture containing 1.5 mM MgCl₂, 0.25 mM dNTPs, 1.5 µM of each primer, reaction buffer and 2.5 U Taq polymerase. The amplification consisted of 94°C for 4 min, 35 cycles of 94°C for 45 s, 55°C

for 45 s (*BsmI*, 58°C for *TaqI* and *ApaI*; 60°C for *FokI*), 72°C for 45 s, and a final extension of 72°C for 5 min.

Ten microlitres of the four PCR fragments generated were subjected to restriction enzyme digestion using *BsmI*, *TaqI*, *ApaI* and *FokI* restriction endonuclease, respectively. The digested PCR products were electrophoresed on a 2% gelose gel and were stained with ethidium bromide to visualize DNA fragments.

Statistical methods

The genotypic distributions of the four SNP sites were all tested for Hardy-Weinberg equilibrium by using Stata 9.0. The distributions of genotypes and allele frequencies in the disease and control groups were compared using the chi-squared test. The risk associated with individual alleles or genotypes was calculated as the odds ratio (OR) with 95% confidence intervals (95% CI). Statistical significance in all tests was determined at $p < 0.05$.

Results

The genotype distributions of all four SNP sites were in accordance with Hardy-Weinberg equilibrium for the control subjects and patients. The PCR-RFLP results showed that no BB or tt genotype was detected in any subject. Genotypic and allelic frequencies involving VDR *BsmI*, *TaqI* and *ApaI* evidenced no differences between the GAgP patients and the control subjects (Table 2).

There was, however, significant difference in the distribution of VDR *FokI* genotypes and the allelic frequencies between the GAgP group and healthy control group. The detected frequency of FF genotype was signifi-

Table 1. The primers for analysing SNPs of the VDR gene (according to GenBank accession no. NT_029419)

SNP	Primers	Fragment
<i>BsmI</i>	5'-ATACCTACTTTGCTGTTTGC-3' 5'-AGCCCATCTCCATTCCTTG-3'	500 bp
<i>TaqI</i> , <i>ApaI</i>	5'-GGAGAAGTCACTGGAGGGC-3' 5'-GGATCATCTTGGCATAGAGC-3'	307 bp
<i>FokI</i>	5'-AGCTGGCCCTGGCACTGACTCTGGCTC-3' 5'-ATGGAAACACCTTGCTTCTTCTCCCTC-3'	267 bp

Table 2. Distribution of VDR genotypes and alleles in GAgP patients and control subjects

SNP	Control subjects (%)	GAgP patients (%)	χ^2	OR
<i>BsmI</i>				
Genotype N	<i>n</i> = 53	<i>n</i> = 51		
BB	0 (0)	0 (0)	—	—
Bb	4 (7, 55)	3 (5, 88)	$\chi^2 = 0.11$	—
bb	49 (92,45)	48 (94 12),	<i>p</i> = 0.73	—
Allele N	<i>n</i> = 106	<i>n</i> = 102		
B	4 (3, 77)	3 (2, 94)	$\chi^2 = 0.11$	—
b	102 (96,23)	99 (97 06),	<i>p</i> = 0.74	—
<i>TaqI</i>				
Genotype N	<i>n</i> = 53	<i>n</i> = 51		
TT	46 (86, 79)	45 (88,24)	—	—
Tt	7 (13,21)	6 (11, 76)	$\chi^2 = 0.05$	—
tt	0 (0)	0 (0)	<i>p</i> = 0.82	—
Allele N	<i>n</i> = 106	<i>n</i> = 102		
T	99 (93,40)	96 (94 12),	$\chi^2 = 0.05$	—
t	7 (6, 60)	6 (5, 88)	<i>p</i> = 0.83	—
<i>ApaI</i>				
Genotype N	<i>n</i> = 53	<i>n</i> = 51		
AA	16 (30 19),	15 (29,41)	—	—
Aa	29 (54,72)	25 (49 02),	$\chi^2 = 0.76$	—
aa	8 (09,15)	11 (21, 57)	<i>p</i> = 0.68	—
Allele N	<i>n</i> = 106	<i>n</i> = 102		
A	61 (57,55)	55 (53,92)	$\chi^2 = 0.28$	—
a	45 (42,45)	47 (46 08),	<i>p</i> = 0.60	—
<i>FokI</i>				
Genotype N	<i>n</i> = 53	<i>n</i> = 51		
FF	9 (16,98)	19 (37,25)	—	FF vs. (Ff + ff)
Ff	27 (50,94)	23 (45 10),	$\chi^2 = 6.32$	OR = 2.90
ff	17 (32 08),	9 (17,65)	<i>p</i> = 0.04	95% CI = 1.16–7.24
Allele N	<i>n</i> = 106	<i>n</i> = 102		F vs. f
F	45 (42,45)	61 (59,80)	$\chi^2 = 6.26$	OR = 2.02
f	61 (57,55)	41 (40 20),	<i>p</i> = 0.01	95% CI = 1.16–3.50

cantly higher in the GAgP group than in the healthy control subjects (37.25 vs. 16.98%, *p* < 0.05). The frequency of F allele was also significantly higher in the GAgP group (59.80 vs. 42.45%, *p* < 0.05). Individuals in the Chinese population with the F allele seemed to be more likely to develop GAgP (FF vs. Ff + ff, OR = 2.90, 95% CI = 1.16–7.24; F vs. f, OR = 2.02, 95% CI = 1.16–3.50; Table 2).

Discussion

Genetic polymorphisms which influence key biological processes may contribute to susceptibility to a variety of diseases, including periodontitis. In particular, the primary clinical feature of GAgP is rapid and severe irreversible alveolar bone loss. There is an increasingly prevalent point of view that periodontitis patients, especially GAgP patients, evidence more clearly

defined genetic factors (6). Candidate genes which have been studied include genes for interleukin (IL)-1, IL-6 receptor, TNF- α , Fc γ R and MMP (7–11).

Genetic polymorphisms, such as SNPs, may influence disease in a complex way, acting with other genetic variants and environmental factors to influence disease susceptibility and progression. Many studies have revealed that certain SNPs may be associated with the susceptibility to aggressive periodontitis in Chinese population, including the IL-1A + 4845 and IL-1B –511 polymorphisms (12), TIMP2 –418 G to C gene polymorphism (11) and MMP-1 promoter region of –1607 bp (13).

In this study, the detected frequency of FF genotype was significantly higher in GAgP patients than in healthy control subjects. The frequency of allele F was also significantly higher in

the GAgP group. No evidence indicated that VDR *BsmI*, *TaqI* or *ApaI* polymorphisms were associated with GAgP in the Chinese population. These results suggest that the FF genotype might be a risk indicator for the susceptibility to GAgP of the Chinese population (OR = 2.90). Carriage of the more frequent allele F of VDR *FokI* gene significantly increased the risk of developing GAgP (OR = 2.02). A study of the Korean population by Park came to a similar conclusion (14). In their study, they found that the VDR *FokI* FF genotype was associated with an increased risk for GAgP (OR = 1.83), but the VDR *BsmI* and *TaqI* polymorphisms were not associated with GAgP. In another study of a Japanese population, they found that VDR *BsmI* polymorphism was not associated with generalized early onset periodontitis (G-EOP) either (15). There were, however, different conclusions in some other studies of Asian aggressive periodontitis (or early onset periodontitis, EOP) patients. In studies by Sun and Tachi, they found that VDR *TaqI* Tt genotype might be a risk indicator for the susceptibility to aggressive periodontitis of Chinese patients (16,17). A possible explanation for these different findings was the observed differences in the disease phenotype. There might also be variations in the frequency of the VDR minor allele in different Asian populations.

The human VDR is a ligand-regulated transcription factor that mediates the actions of the 1,25-dihydroxyvitamin D₃ hormone to effect bone mineral homeostasis. The VDR mediates the hormonal function of vitamin D and regulates a variety of downstream functions. Vitamin D receptor *BsmI* and *ApaI* polymorphisms in this study are located at intron 8 and VDR *TaqI* polymorphism is located at exon 9, but they are synonymous mutations. None of the three SNPs can lead to a change of the protein structure. This might be the key reason why there was no association between the VDR *BsmI*, *ApaI* and *TaqI* SNPs and GAgP. In addition, the VDR gene contains two potential translation initiation (ATG) sites. The *FokI* polymorphism, which

occurs at the first start codon in exon 2, changes the nucleotide sequence to ACG (18). Alleles with this polymorphism initiate translation three codons downstream, resulting in a protein (424 amino acids, the F allele) that is three amino acids shorter than wild type. The short 424-amino-acid VDR interacted more efficiently transactivates with the transcription factor TFII B than the 427-amino-acid VDR *in vitro*, and the shorter VDR also transactivates vitamin D target genes more efficiently. The shorter VDR transmits stronger bone resorption and inflammation signals (19,20). We consider that is the main reason why VDR FF genotype is associated with an increased risk of GAgP.

References

- Hassell TM, Harris EL. Genetic influences in caries and periodontal diseases. *Crit Rev Oral Biol Med* 1995;**6**:319–342.
- Lin NU, Malloy PJ, Sakati N, al-Ashwal A, Feldman D. A novel mutation in the deoxyribonucleic acid-binding domain of the vitamin D receptor causes hereditary 1,25-dihydroxyvitamin D-resistant rickets. *J Clin Endocrinol Metab* 1996;**81**:2564–2569.
- Malloy PJ, Eccleshall TR, Gross C, Van Maldergem L, Bouillon R, Feldman D. Hereditary vitamin D resistant rickets caused by a novel mutation in the vitamin D receptor that results in decreased affinity for hormone and cellular hyporesponsiveness. *J Clin Invest* 1997;**99**:297–304.
- Selvaraj P, Chandra G, Jawahar MS, Rani MV, Rajeshwari DN, Narayanan PR. Regulatory role of vitamin D receptor gene variants of *BsmI*, *ApaI*, *TaqI*, and *FokI* polymorphisms on macrophage phagocytosis and lymphoproliferative response to *Mycobacterium tuberculosis* antigen in pulmonary tuberculosis. *J Clin Immunol* 2004;**24**:523–532.
- Armitage GC. Development of a classification system for periodontal diseases and conditions. *Ann Periodontol* 1999;**4**:1–6.
- Hodge P, Michalowicz B. Genetic predisposition to periodontitis in children and young adults. *Periodontol* 2000 2001;**26**: 113–134.
- Quappe L, Jara L, Lopez NJ. Association of interleukin-1 polymorphisms with aggressive periodontitis. *J Periodontol* 2004;**75**:1509–1515.
- Galicía JC, Tai H, Komatsu Y, Shimada Y, Ikezawa I, Yoshie H. Interleukin-6 receptor gene polymorphisms and periodontitis in a non-smoking Japanese population. *J Clin Periodontol* 2006;**33**: 704–709.
- Trevilatto PC, Tramontina VA, Machado MA, Goncalves RB, Sallum AW, Line SR. Clinical, genetic and microbiological findings in a Brazilian family with aggressive periodontitis. *J Clin Periodontol* 2002;**29**:233–239.
- de Souza RC, Colombo AP. Distribution of FcγRIIa and FcγRIIb genotypes in patients with generalized aggressive periodontitis. *J Periodontol* 2006;**77**: 1120–1128.
- Chen D, Wang Q, Ma ZW *et al*. MMP-2, MMP-9 and TIMP-2 gene polymorphisms in Chinese patients with generalized aggressive periodontitis. *J Clin Periodontol* 2007;**34**:384–389.
- Li QY, Zhao HS, Meng HX *et al*. Association analysis between interleukin-1 family polymorphisms and generalized aggressive periodontitis in a Chinese population. *J Periodontol* 2004;**75**:1627–1635.
- Cao Z, Li C, Jin L, Corbet EF. Association of matrix metalloproteinase-1 promoter polymorphism with generalized aggressive periodontitis in a Chinese population. *J Periodont Res* 2005;**40**:427–431.
- Park KS, Nam JH, Choi J. The short vitamin D receptor is associated with increased risk for generalized aggressive periodontitis. *J Clin Periodontol* 2006;**33**:524–528.
- Yoshihara A, Sugita N, Yamamoto K, Kobayashi T, Miyazaki H, Yoshi H. Analysis of vitamin D and Fcγ receptor polymorphisms in Japanese patients with generalized early-onset periodontitis. *J Dent Res* 2001;**80**:2051–2054.
- Sun JL, Meng HX, Cao CF *et al*. Relationship between vitamin D receptor gene polymorphism and periodontitis. *J Periodont Res* 2002;**37**:263–267.
- Tachi Y, Shimpuku H, Nosaka Y *et al*. Association of vitamin D receptor gene polymorphism with periodontal diseases in Japanese and Chinese. *Nucl Acids Res Suppl* 2001;**1**:111–112.
- Saijo T, Ito M, Takeda E *et al*. A unique mutation in the vitamin D receptor gene in three Japanese patients with vitamin D-dependent rickets type II: utility of single-strand conformation polymorphism analysis for heterozygous carrier detection. *Am J Hum Genet* 1991;**49**:668–673.
- Jurutka PW, Remus LS, Whitfield GK *et al*. The polymorphic N terminus in human vitamin D receptor isoforms influences transcriptional activity by modulating interaction with transcription factor IIB. *Mol Endocrinol* 2000;**14**: 401–420.
- Whitfield GK, Remus LS, Jurutka PW *et al*. Functionally relevant polymorphisms in the human nuclear vitamin D receptor gene. *Mol Cell Endocrinol* 2001;**177**:145–159.

CIMS Papers' World

Here scientific papers on Computer Integrated Manufacturing System(CIMS) and other fields!for you FREELY!

<http://www.cimspaper.com>